

*Weight control strategies that are both safe and effective are needed to reduce the rate of the current obesity epidemic. In a recent study, people incorporating fortified meal replacements were more likely to have adequate essential nutrient intakes compared to a group following a more traditional food group diet.*

Dietary supplements and fortified meal replacements help ensure nutritional adequacy during energy-restricted diets for weight loss

Weight control strategies that are both safe and effective are needed to reduce the rate of the current obesity epidemic. A recent study compared the macronutrient and micronutrient levels in the foods chosen by women following two different weight reduction programs.

Ninety-six generally healthy overweight or obese women randomly placed into two treatment groups: Traditional Food Group (TFG) or a Meal Replacement Group (MRG). The MRG included the use of 1-2 meal replacement drinks or bars per day. Both groups aimed to restrict energy levels to approximately 1,300 calories per day.

After one year, weight loss was not significantly different between the groups, and both groups had macronutrient (Carbohydrate:Protein:Fat) ratios that were within the ranges recommended. Both groups experienced an improved dietary pattern with respect to decreased saturated fat, cholesterol, and sodium, with increased total servings/day of fruits and vegetables. However, the TFG had a significantly lower dietary intake of several vitamins and minerals compared to the MRG and was at greater risk for inadequate intake.

Although both groups successfully lost weight while improving overall dietary adequacy, the group incorporating fortified meal replacements tended to have a more adequate essential nutrient intake compared to the group following a more traditional food group diet. This study supports the need to incorporate fortified foods and/or dietary supplements while following an energy-restricted diet for weight loss.

*Nutr J. 2007 Jun 25;6:12*